

Reading the Results

Evaluation

Colonies of *T. equigenitalis* are small (2-3 mm), smooth with complete edges, glossy and yellowish gray.²

Limitations

For Veterinary and Research Use Only

Plates should be examined for contaminants after the first 24 hours of incubation. Laboratories should be aware that certain countries and/or states may require the prolonged incubation periods of specific confirmation techniques as standard procedures and should therefore ascertain the particular local or regional requirements for specific testing and reporting and/or indicate the specific isolation and testing methods used for their cultural findings. Definite confirmation of any species requires a range of staining, biochemical testing, antibody agglutination or immunofluorescent testing.

Symbol glossary: biomeddiagnostics.com/l/symbol-glossary

References:

1. Pelczar and Vera (1949) Milk Plant Monthly 38:30.
2. World Organization for Animal Health (OIE) (2019). Manual of Diagnostic Tests and Vaccines for Terrestrial Animals (Terrestrial Manual). oie.int/standard-setting/terrestrial-manual/access-online.

Document Revision History

Rev. C, October 2019

New format; added new catalog numbers, document revision history; specified 18-25°C instead of room temperature; reorganized and retitled some sections

Scan for additional product information



Petri Chocolate Eugon w/ 10% Horse Blood

| | | |
|-------------|------------|-------------------|
| Catalog No. | 12-134-001 | 10 pack, unsealed |
| Catalog No. | 12-134-002 | 10 pack, sealed |

Not available in all countries; please inquire.

For Veterinary and Research Use Only

Manufactured by:
Biomed Diagnostics, Inc.
1388 Antelope Road
White City, OR 97503 USA
biomeddiagnostics.com

Download



Certificate
of Analysis

Introduction

Intended Use

Chocolate Eugin Agar with 10% Horse Blood (CEA-HB) is a non-selective medium to aid in the culture based isolation and identification of a wide variety of fastidious pathogenic bacteria.

Description and Principle

Our Eugin base is prepared according to the formula described by Pelczar and Vera, which was developed to obtain eugenic (luxuriant) growth of fastidious microorganisms.¹ The CEA-HB medium, enriched with 10% “chocolate” horse blood (heat-lysed), supports the growth of pathogenic fungi, fastidious *Neisseria*, *Brucella*, *Taylorella*, *Lactobacillus*, *Streptococcus*, *Haemophilus* and others. CEA-HB is often used as a non-selective isolation medium for Contagious Equine Metritis (CEM) screening. CEM is an inflammation of the endometrium of mares caused by *T. equigenitalis*, which usually results in temporary infertility. CEA-HB is suitable for the direct plating of fresh swabs of properly controlled samples transported in Amie’s or other applicable transport media. For a selective medium for CEM, try our Timoney’s CEM Agar (Cat. Nos. 12-144-001, 12-144-002).²

Reagents and Appearance

CEA-HB medium appears chocolate brown and contains agar, peptone nutrients, horse blood, dextrose, L-cysteine, sodium sulfite and salt with a final pH of 7.0 ± 0.1 at 25°C.

Precautions, Safety and Disposal

Read the Safety Data Sheets (SDSs) and follow the handling instructions. Wear appropriate protective eyewear, clothing and gloves.

Once the petri has been inoculated, re-open only in a biological safety cabinet. Because of the potential for containing infectious materials, the petri must be destroyed by autoclaving at 121°C for 20 minutes.

Storage

Upon receipt, store CEA-HB Agar under refrigeration (2-8°C). Medium can be kept for one day at ambient temperature. Avoid freezing or prolonged storage at temperatures above 40°C. Petri dishes should be received and stored upside down to minimize condensation on the surface of the agar. Do not open until ready to use. Do not use if the medium shows signs of deterioration or contamination.

Shelf Life


Petri Chocolate Eugin agar expires 6 months from date of manufacture.

Procedure

Materials Provided

- Petri Chocolate Eugin w/ 10% Horse Blood

1



Allow the petri to warm to 18-25°C.

Lift the inverted petri base off the lid.

For sealed petri dishes pull off the seal with care while in the inverted position and discard in regular trash.

Incubation

Incubate **inverted** at 30-37°C for 18-72 hours (up to 14 days) in 5% to 7% CO₂ environment with a 85% to 98% relative humidity range.

Quality Control


This product has been tested and meets the CLSI (formerly NCCLS) Approved Standard for commercially prepared media (M22-A3). At the time of manufacture, quality control testing is performed on each lot of the Petri Chocolate Eugin w/ 10% Horse Blood. The ability of the media to support growth and demonstrate expected biochemical reactions and morphology is verified by lot.

All Biomed product lots are performance verified with ATCC® microbe strains. Product performance is also verified periodically throughout the marked shelf life of each lot.

Materials Required but Not Provided

- Sterile inoculating tool (e.g, cotton swab/forceps/scalpel blade)
- Laboratory incubator capable of incubation at 30-37°C in 5% to 7% CO₂ environment with a 85% to 98% relative humidity range

2



Streak the sample onto the agar surface.

Immediately label the petri with subject and sample information and date.

Organisms used for quality control testing

| Organism | ATCC | Colony Appearance |
|-------------------------|-------|--|
| <i>T. equigenitalis</i> | 35865 | Small, smooth, yellowish grey, cytochrome-oxidase positive |
| <i>E. coli</i> | 25922 | Growth, cytochrome-oxidase negative |
| <i>S. xylosus</i> | 29971 | Growth, cytochrome-oxidase negative |
| <i>C. albicans</i> | 90028 | Growth cytochromeoxidase negative |

(Terrestrial Manual)²